

Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed ☐ for     
☒ Limits Preview/Index History Clipboard Details

About Entrez

Abstract    Text

Text Version

☐ 1: Ann Hum Genet. 1999 Jul;63 ( Pt 4):277-84.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities



## Allelic diversity of the human plasma alpha(1,3)fucosyltransferase gene (FUT6).

Pang H, Koda Y, Soejima M, Schlaphoff T, du Toit ED, Kimura H.

Department of Forensic Medicine, Kurume University School of Medicine, Fukuoka, Japan.

The 1080-bp coding region of the human plasma alpha(1,3)fucosyltransferase gene (FUT6) was sequenced in a total of 161 individuals (322 chromosomes) drawn from three populations, involving 56 Africans (Xhosa), 52 European-Africans of South Africa, and Japanese. In addition to six reported base substitutions, eleven new base substitutions and a single base insertion were found in the coding region of the FUT6. Eleven functional and four null alleles were encountered, of which 10 alleles were novel alleles identified in this study. Two null alleles have been identified previously, whereas two novel null alleles, which contained a single base (cytosine) insertion at nucleotide 499, were found in a Xhosa population. The allelic distributions of FUT6 were different among these three populations. The heterozygosity of FUT6 was 0.860, 0.699, and 0.632, in Xhosa, European-African (South Africa), and in Japanese populations, respectively. The extensive DNA sequence diversity of the FUT6 may be suitable for application as a tool in genetic studies for modern human evolution.

PMID: 10738539 [PubMed - indexed for MEDLINE]


Abstract    Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [ ] Go Clear  
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Biochim Biophys Acta. 1999 Dec 6;1473(1):9-20.

Related Articles, L

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-UtilitiesPubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
CubbyRelated Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

## The alpha1-6-fucosyltransferase gene and its biological significance.

Miyoshi E, Noda K, Yamaguchi Y, Inoue S, Ikeda Y, Wang W, Ko JH, Uozumi N, L W, Taniguchi N.

Department of Biochemistry, Osaka University Medical School, Room B-1, 2-2 Yamadaoka, Suita, Osaka, Japan.


GDP-L-Fuc:N-acetyl-beta-D-glucosaminide alpha1-6-fucosyltransferase (alpha1-6FucT catalyzes the transfer of fucose from GDP-Fuc to N-linked type complex glycoproteins. This enzyme was purified from a human fibroblast cell line, porcine brain, a human gast cancer cell line and human blood platelets. cDNA cloning of porcine and human alpha1-6FucT was performed from a porcine brain and gastric cancer cell cDNA libraries, respectively. Their homology is 92.2% at the nucleotide level and 95.7% at the amino ac level. No putative N-glycosylation sites were found in the predicted amino acid sequence. No homology to other fucosyltransferases such as alpha1-2FucT, alpha1-3FucT and alpha1-4FucT was found except for a region consisting of nine amino acids. The alpha1-6FucT gene is located at chromosome 14q24.3, which is also a different location from other fucosyltransferases reported to date. The alpha1-6FucT gene is the oldest gene fam in the phylogenic trees among the nine cloned fucosyltransferase genes. alpha1-6FucT is widely expressed in various rat tissues and the expression of alpha1-6FucT in the liver is enhanced during hepatocarcinogenesis of LEC rats which develop hereditary hepatitis an hepatomas. In cases of human liver diseases, alpha1-6FucT is expressed in both hepatom tissues and their surrounding tissues with chronic liver disease, but not in the case of normal liver. Serum alpha1-6-fucosylated alpha-fetoprotein (AFP) has been employed fo an early diagnosis of patients with hepatoma. The mechanisms by which alpha1-6 fucosylation of AFP occurs in the hepatoma is not due to the up-regulation of alpha1-6FucT alone. Interestingly, when the alpha1-6FucT gene is transfected into Hep3B, a human hepatoma cell line, tumor formation in the liver of nude mice after splenic injecti is dramatically suppressed. In this review, we focus on alpha1-6FucT and summarize its properties, gene expression and biological significance.

### Publication Types:

- Review
- Review, Tutorial

PMID: 10580126 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed for [ ] Go Clear  
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

1: J Immunol. 1999 Oct 1;163(7):3746-52.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

FREE full text article at  
[www.jimmunol.org](http://www.jimmunol.org)

## Alpha(1,3)-fucosyltransferase VII and alpha(2,3)-sialyltransferase IV are up-regulated in activated CD4 T cells and maintained after their differentiation into Th1 and migration into inflammatory sites.

Blander JM, Visintin I, Janeway CA Jr, Medzhitov R.


Section of Immunobiology, Yale University School of Medicine, Howard Hughes Medical Institute, New Haven, CT 06520, USA.

Activated Th1 CD4 T cells bind to P-selectin and migrate into inflamed tissue, whereas Th2 cells do not. We show that alpha(1, 3)-fucosyltransferase VII (FucT-VII) and alpha(2, 3)-sialyltransferase IV (ST3GalIV), which are crucial for the biosynthesis of functional P selectin ligands, are absent in naive CD4 T cells, but are rapidly up-regulated upon activation. Th1 or Th2 differentiation in the presence of polarizing cytokines leads to down-regulation of FucT-VII mRNA selectively in Th2 but not in Th1 cells. Influencing the differentiation by varying the priming dose of antigenic peptide results in similar FucT-VII down-regulation only in Ag-specific Th2 cells. ST3GalIV levels remain elevated. FucT-VII and ST3GalIV mRNAs are also up-regulated by Th1 cells primed in vivo and recruited into the lymph nodes draining delayed-type hypersensitivity sites. We identify FucT-VII gene expression as a principal difference between Th1 and Th2 cells, and underscore the importance of FucT-VII and ST3GalIV expression for the biosynthesis of functional selectin ligands.

PMID: 10490970 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
Freedom of Information Act | Disclaimer



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for #1 AND #6 Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

Field: Title, Limits: Publication Date from 1980 to 1999

Display Summary Show: 20 Sort Send to Text

Items 1-14 of 14

One page

- ☐ 1: [Leiter H, Mucha J, Staudacher E, Grimm R, Glossl J, Altmann F.](#) Related Articles, Li  
Purification, cDNA cloning, and expression of GDP-L-Fuc:Asn-linked GlcNAc alpha1, fucosyltransferase from mung beans.  
J Biol Chem. 1999 Jul 30;274(31):21830-9.  
PMID: 10419500 [PubMed - indexed for MEDLINE]
- ☐ 2: [Kimura N, Mitsuoka C, Kanamori A, Hiraiwa N, Uchimura K, Muramatsu T, Tamatani T, Kansas GS, Kannagi R.](#) Related Articles, Li  
Reconstitution of functional L-selectin ligands on a cultured human endothelial cell line by cotransfection of alpha1-->3 fucosyltransferase VII and newly cloned GlcNAc beta:6 sulfotransferase cDNA.  
Proc Natl Acad Sci U S A. 1999 Apr 13;96(8):4530-5.  
PMID: 10200296 [PubMed - indexed for MEDLINE]
- ☐ 3: [Hallouin F, Goupille C, Bureau V, Meflah K, Le Pendu J.](#) Related Articles, Li  
Increased tumorigenicity of rat colon carcinoma cells after alpha1,2-fucosyltransferase FTA anti-sense cDNA transfection.  
Int J Cancer. 1999 Feb 9;80(4):606-11.  
PMID: 9935164 [PubMed - indexed for MEDLINE]
- ☐ 4: [Sepp A, Skacel P, Lindstedt R, Lechler RI.](#) Related Articles, Li  
Expression of alpha-1,3-galactose and other type 2 oligosaccharide structures in a porcine endothelial cell line transfected with human alpha-1,2-fucosyltransferase cDNA.  
J Biol Chem. 1997 Sep 12;272(37):23104-10.  
PMID: 9287311 [PubMed - indexed for MEDLINE]
- ☐ 5: [Yanagidani S, Uozumi N, Ihara Y, Miyoshi E, Yamaguchi N, Taniguchi N.](#) Related Articles, Li  
Purification and cDNA cloning of GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha1-6 fucosyltransferase (alpha1-6 FucT) from human gastric cancer MKN45 cells.  
J Biochem (Tokyo). 1997 Mar;121(3):626-32.  
PMID: 9133635 [PubMed - indexed for MEDLINE]
- ☐ 6: [Gorelik E, Xu F, Henion T, Anaraki F, Galili U.](#) Related Articles, Li  
Reduction of metastatic properties of BL6 melanoma cells expressing terminal fucose (alpha)1-2-galactose after alpha1,2-fucosyltransferase cDNA transfection.  
Cancer Res. 1997 Jan 15;57(2):332-6.  
PMID: 9000578 [PubMed - indexed for MEDLINE]
- ☐ 7: [Hiraiwa N, Dohi T, Kawakami-Kimura N, Yumen M, Ohmori K, Maeda M, Kannagi R.](#) Related Articles, Li  
Suppression of sialyl Lewis X expression and E-selectin-mediated cell adhesion in cultured human lymphoid cells by transfection of antisense cDNA of an alpha1-->3 fucosyltransferase (Fuc-T VII).  
J Biol Chem. 1996 Dec 6;271(49):31556-61.  
PMID: 8940172 [PubMed - indexed for MEDLINE]

☐ **8:** [Natsuka S, Gersten KM, Zenita K, Kannagi R, Lowe JB.](#)

[Related Articles, Li](#)



Molecular cloning of a cDNA encoding a novel human leukocyte alpha-1,3-fucosyltransferase capable of synthesizing the sialyl Lewis x determinant.

J Biol Chem. 1994 Aug 12;269(32):20806. No abstract available.

PMID: 8051184 [PubMed - indexed for MEDLINE]

☐ **9:** [Natsuka S, Gersten KM, Zenita K, Kannagi R, Lowe JB.](#)

[Related Articles, Li](#)



Molecular cloning of a cDNA encoding a novel human leukocyte alpha-1,3-fucosyltransferase capable of synthesizing the sialyl Lewis x determinant.

J Biol Chem. 1994 Jun 17;269(24):16789-94. Erratum in: J Biol Chem 1994 Aug 12;269(32):20806.

PMID: 8207002 [PubMed - indexed for MEDLINE]

☐ **10:** [Lowe JB, Stoolman LM, Nair RP, Larsen RD, Behrend TL, Marks RM.](#)

[Related Articles, Li](#)



A transfected human fucosyltransferase cDNA determines biosynthesis of oligosaccharide ligand(s) for endothelial-leukocyte adhesion molecule I.

Biochem Soc Trans. 1991 Aug;19(3):649-53. No abstract available.

PMID: 1723704 [PubMed - indexed for MEDLINE]

☐ **11:** [Lowe JB, Stoolman LM, Nair RP, Larsen RD, Berhend TL, Marks RM.](#)

[Related Articles, Li](#)



ELAM-1--dependent cell adhesion to vascular endothelium determined by a transfected human fucosyltransferase cDNA.

Cell. 1990 Nov 2;63(3):475-84.

PMID: 1699667 [PubMed - indexed for MEDLINE]

☐ **12:** [Larsen RD, Ernst LK, Nair RP, Lowe JB.](#)

[Related Articles, Li](#)



Molecular cloning, sequence, and expression of a human GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase cDNA that can form the H blood group antigen.

Proc Natl Acad Sci U S A. 1990 Sep;87(17):6674-8.

PMID: 2118655 [PubMed - indexed for MEDLINE]

☐ **13:** [Kukowska-Latallo JF, Larsen RD, Nair RP, Lowe JB.](#)

[Related Articles, Li](#)



A cloned human cDNA determines expression of a mouse stage-specific embryonic antigen and the Lewis blood group alpha(1,3/1,4)fucosyltransferase.

Genes Dev. 1990 Aug;4(8):1288-303.

PMID: 1977660 [PubMed - indexed for MEDLINE]

☐ **14:** [Prieels JP, Monnom D, Perraudin JP, Finne J, Burger M.](#)

[Related Articles, Li](#)



Enzymic properties of an N-acetylglucosaminide 3-alpha-L-fucosyltransferase of a wheat-germ agglutinin-resistant melanoma clone.

Eur J Biochem. 1983 Feb 1;130(2):347-51.

PMID: 6687456 [PubMed - indexed for MEDLINE]

Display  Show:  Sort

Items 1-14 of 14

One page

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed for [ ] Go Clear  
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Biol Chem. 1994 Aug 12;269(32):20806.

Related Articles, L

FREE full text article at  
[www.jbc.org](http://www.jbc.org)

**Molecular cloning of a cDNA encoding a novel human leukocyte alpha-1,3 fucosyltransferase capable of synthesizing the sialyl Lewis x determinant.**

Natsuka S, Gersten KM, Zenita K, Kannagi R, Lowe JB.

PMID: 8051184 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jun 1 2004 06.



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed for Go Clear  
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: J Biol Chem. 1999 Jul 30;274(31):21830-9.

Related Articles, L

FREE full text article at  
[www.jbc.org](http://www.jbc.org)

## Purification, cDNA cloning, and expression of GDP-L-Fuc:Asn-linked GlcNAc alpha1,3-fucosyltransferase from mung beans.


Leiter H, Mucha J, Staudacher E, Grimm R, Glossl J, Altmann F.

Institute of Chemistry, Universitat fur Bodenkultur, Muthgasse 18, 1190 Wien, Austria.  
[faltmann@edv2.boku.ac.at](mailto:faltmann@edv2.boku.ac.at)

Substitution of the asparagine-linked GlcNAc by alpha1,3-linked fucose is a widespread feature of plant as well as of insect glycoproteins, which renders the N-glycan immunogenic. We have purified from mung bean seedlings the GDP-L-Fuc:Asn-linked GlcNAc alpha1,3-fucosyltransferase (core alpha1,3-fucosyltransferase) that is responsible for the synthesis of this linkage. The major isoform had an apparent mass of 54 kDa and isoelectric points ranging from 6.8 to 8.2. From that protein, four tryptic peptides were isolated and sequenced. Based on an approach involving reverse transcriptase-polymerase chain reaction with degenerate primers and rapid amplification of cDNA ends, core alpha1,3-fucosyltransferase cDNA was cloned from mung bean mRNA. The 2200-base pair cDNA contained an open reading frame of 1530 base pairs that encoded a 510-amino acid protein with a predicted molecular mass of 56.8 kDa. Analysis of cDNA derived from genomic DNA revealed the presence of three introns within the open reading frame. Remarkably, from the four exons, only exon II exhibited significant homology to animal and bacterial alpha1,3/4-fucosyltransferases which, though, are responsible for the biosynthesis of Lewis determinants. The recombinant fucosyltransferase was expressed in Sf21 insect cells using a baculovirus vector. The enzyme acted on glycopeptides having glycan structures GlcNAc beta1-2Man alpha1-3(GlcNAc beta1-2Man alpha1-6)Man beta1-4GlcNAc beta1-4GlcNAc beta1-Asn, GlcNAc beta1-2Man alpha1-3(GlcNAc beta1-2Man alpha1-6)Man beta1-4GlcNAc beta1-4(Fuc alpha1-6)GlcNAc beta1-Asn, and GlcNAc beta1-2Man alpha1-3[Man alpha1-3(Man alpha1-6)Man alpha1-6]Man beta1-4GlcNAc beta1-4GlcNAc beta1-Asn but not on, e.g. N-acetylglucosamine. The structure of the core alpha1,3-fucosylated product was verified by high performance liquid chromatography of the pyridylaminated glycan and by its insensitivity to N-glycosidase as revealed by matrix-assisted laser desorption/ionization time of flight mass spectrometry.

PMID: 10419500 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed ☐ for     
☒ Limits Preview/Index History Clipboard Details

About Entrez

Abstract

Text Version

☐ 1: J Biochem (Tokyo). 1997 Mar;121(3):626-32.

[Related Articles, L](#)

Entrez PubMed

Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities

PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

[Privacy Policy](#)

## Purification and cDNA cloning of GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha1-6 fucosyltransferase (alpha1-6 FucT) from human gastric cancer MKN45 cells.

Yanagidani S, Uozumi N, Ihara Y, Miyoshi E, Yamaguchi N, Taniguchi N.

Department of Biochemistry, Osaka University Medical School, Suita.


GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha1-6 fucosyltransferase (alpha1-6 FucT) which catalyzes the transfer of fucose from GDP-Fuc to N-linked type complex glycopeptides, was purified from a culture supernatant of human gastric cancer cell line MKN45. The purification procedures included chromatographies on Q-Sepharose Fast Flow, synthetic GDP-hexanolamine-Sepharose, and GnGn-bi-Asn-Sepharose columns. SDS-PAGE of the purified enzyme gave a major band corresponding to an apparent molecular mass of 60 kDa. The enzyme was recovered in a 12% final yield with an approximately 4,600-fold increase in specific activity. The pH optimum was 7.5, and the enzyme was fully active in the presence of 5 mM EDTA and did not require divalent cations, Mg<sup>2+</sup> and Ca<sup>2+</sup>. Oligonucleotide primers designed from partial amino acid sequences were used to amplify and clone alpha1-6 FucT cDNA from a cDNA library of MKN45 cells. The cDNA encodes 575 amino acids in length, and contains the predicted terminal and internal amino acid sequences derived on lysyl endopeptidase digestion. The homology to porcine brain alpha1-6 FucT is 92.2% at the nucleotide level and 95.7% at amino acid level. No putative N-glycosylation sites were found in the predicted amino acid sequence of the human MKN45 cell enzyme or that of porcine brain. Thus, the enzyme is distinct from other fucosyltransferases which catalyze alpha1-2, alpha1-3, and alpha1-4 fucose addition.

PMID: 9133635 [PubMed - indexed for MEDLINE]

Abstract

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
Department of Health & Human Services  
[Freedom of Information Act](#) | [Disclaimer](#)





Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book  
Search PubMed for [ ] Go Clear  
☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Proc Natl Acad Sci U S A. 1990 Sep;87(17):6674-8.

Related Articles, L

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

FREE full text article at  
www.pnas.org

FREE full text article  
in PubMed Central

## Molecular cloning, sequence, and expression of a human GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase cDNA that can for the H blood group antigen.

Larsen RD, Ernst LK, Nair RP, Lowe JB.

Howard Hughes Medical Institute, Ann Arbor, MI.

We have previously used a gene-transfer scheme to isolate a human genomic DNA fragment that determines expression of a GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase [alpha(1,2)FT; EC 2.4.1.69]. Although this fragment determined expression of an alpha(1,2)FT whose kinetic properties mirror those of the human H blood group alpha(1,2)FT, their precise nature remained undefined. We describe here the molecular cloning, sequence, and expression of a human cDNA corresponding to these human genomic sequences. When expressed in COS-1 cells, this cDNA directs expression of cell surface H structures and a cognate alpha(1,2)FT activity with properties analogous to the human H blood group alpha(1,2)FT. The cDNA sequence predicts a 365-amino acid polypeptide characteristic of a type II transmembrane glycoprotein with a domain structure analogous to that of other glycosyltransferases but without significant primary sequence similarity to these or other known proteins. To directly demonstrate that the cDNA encodes an alpha(1,2)FT, the COOH-terminal domain predicted to be Golgi-resident was expressed in COS-1 cells as a catalytically active, secreted, and soluble protein A fusion peptide. Southern blot analysis showed that this cDNA identifies DNA sequences syntenic to the human H locus on chromosome 19. These results strongly suggest that this cloned alpha(1,2)FT cDNA represents the product of the human H blood group locus.

PMID: 2118655 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
Freedom of Information Act | Disclaimer